



User name: \_\_\_\_\_  
Password: \_\_\_\_\_   
[Register](#) [Forgotten Password](#) [Athens Login](#)

[Synergy Home](#) | [Browse](#) | [Search](#) | [My Synergy](#) | [Books Online](#) | [Resources](#) | [About](#) | [Help](#)



## JOURNAL OF Veterinary Pharmacology and Therapeutics

### Journal Menu

[Table of Contents](#)  
[List of Issues](#)

### Tools

[Email this article](#)  
[Add to favorite articles](#)  
[Export this citation](#)  
[Alert me when this article is cited: Email | RSS](#)  
(What is this?)

[View PubMed citation](#)  
[View ISI citation](#)

### Publication history

**Issue online:**  
05 Jan 2002

[Home](#) > [List of Issues](#) > [Table of Contents](#) > [Article Abstract](#)


## Journal of Veterinary Pharmacology and Therapeutics

Volume 21 Issue 1 Page 74 - February 1998

**To cite this article:** ELLIOTT, BRADY (1998)

Characterization of vasodilatory adenosine receptors in equine digital veins

Journal of Veterinary Pharmacology and Therapeutics 21 (1), 74-81.  
doi:10.1046/j.1365-2885.1998.00106.x



[Prev Article](#) | [Next Article](#)

### Original Article

## Characterization of vasodilatory adenosine receptors in equine digital veins

ELLIOTT & BRADY

<sup>1</sup>Department of Veterinary Basic Sciences, Royal Veterinary College,  
Royal College Street, London NW1 0TU, UK

*Correspondence to:* Elliott Department of Veterinary Basic  
Sciences, Royal Veterinary College, Royal College Street,  
London NW1 0TU, UK

### Abstract

Isolated equine digital veins (EDVs) which had been denuded of their endothelium were used to study adenosine receptors causing vasodilation. When the blood vessel wall tension was raised with the thromboxanemimetic, U44069 (30 nM), the order of vasodilator potency of adenosine receptor agonists was: 5'-N-ethylcarboxamidoadenosine (NECA) > 2-*p*-(2-carboxyethyl)phenyl amino-5'-N-ethylcarboxamido-adenosine (CGS 21680) > 5'-N-methylcarboxamido-adenosine (MECA) >> N<sup>6</sup>-cyclohexyladenosine (CHA) > N<sup>6</sup>-cyclopentyladenosine (CPA) > N<sup>6</sup>-2-(4-Aminophenyl) ethyladenosine (APNEA) > adenosine. Removal of the endothelium had no significant effect on the responses to NECA. The adenosine receptor antagonists, 8-cyclopentyl-1,3-dipropylxanthine (DPCPX; A<sub>1</sub>-selective) and xanthine amine cogener (XAC; non-selective

*J Neurophysiol* 79: 501-510, 1998;  
0022-3077/98 \$5.00

The Journal of Neurophysiology Vol. 79 No. 2 February 1998, pp. 501-510  
Copyright ©1998 by the American Physiological Society

## Adenosine A1 Receptors Mediate Retinotectal Presynaptic Inhibition: Uncoupling by C-Kinase and Role in LTP During Regeneration

Chunyi Zhang and John T. Schmidt

Department of Biological Sciences and Neurobiology Research Center,  
State University of New York, Albany, New York 12222

### ► ABSTRACT

**Zhang, Chunyi and John T. Schmidt.** Adenosine A1 receptors mediate retinotectal presynaptic inhibition: uncoupling by C-kinase and role in LTP during regeneration. *J. Neurophysiol.* 79: 501-510, 1998. Presynaptic adenosine receptors inhibit transmitter release at many synapses and are known to

exist on retinotectal terminals. In this paper we show that adenosine decreases retinotectal field potentials by ~30% and investigate the mechanism. First, as judged by the effects of specific calcium channel blockers, retinotectal transmission was mediated almost exclusively by N-type calcium channels, which are known to be modulated by adenosine A1 receptors. Transmission was completely blocked by either  $\omega$ -Conotoxin GVIA (–100%, N-type blocker) or  $\omega$ -Conotoxin MVIIC (–99%, N-, P- and Q-type blocker) and was not significantly affected by  $\omega$ -Agatoxin IVA [ $+1.7 \pm 9.3\%$  (SE), P-, Q-type blocker], but was augmented slightly by nifedipine ( $+9.3 \pm 2.1\%$ , L-type blocker). Second, the adenosine inhibition was presynaptic, as indicated by a 43% increase in paired-pulse facilitation. Third, the selective A1 agonist cyclohexyl adenosine (CHA) at 50 nM caused a 21% decrease in amplitude and the selective A2 agonist *N*<sup>6</sup>-[2-(3,5-dimethoxyphenyl)-2-(2-methylphenyl)-ethyl]adenosine (DPMA) at 100 nM caused a 24% increase. Fourth, the selective A1 antagonist 8-cyclopentyl-1,3-dipropylxanthine (DPCPX) alone produced an increase in the field potential, suggesting a tonic inhibition mediated by endogenous adenosine. Fifth, pertussis toxin eliminated adenosine inhibition implicating G<sub>i</sub> or G<sub>o</sub> protein coupling. Sixth, C-kinase activation eliminated the A1-mediated inhibition. In regenerating projections, adenosine also caused a decrease in transmission ( $-30 \pm 12\%$ ), but after induction of long-term potentiation (LTP) via trains of stimuli or via treatment with the phosphatase inhibitor okadaic acid, the adenosine response was converted to an augmentation. Because LTP is associated with C-kinase activation, this is consistent with C-kinase uncoupling the A1 receptor from inhibiting

- Abstract
- ▼ Introduction
- ▼ Methods
- ▼ Results
- ▼ Discussion
- ▼ References

### This Article

- **Abstract** FREE
- **Full Text (PDF)**
- **Alert me when this article is cited**
- **Alert me if a correction is posted**
- **Citation Map**

### Services

- **Email this article to a friend**
- **Similar articles in this journal**
- **Similar articles in PubMed**
- **Alert me to new issues of the journal**
- **Download to citation manager**
- **Cited by other online articles**

### Google Scholar

- **Articles by Zhang, C.**
- **Articles by Schmidt, J. T.**
- **Articles citing this Article**

### PubMed

- **PubMed Citation**
- **Articles by Zhang, C.**
- **Articles by Schmidt, J. T.**



Content Types Subjects

English



Athens Authentication Point

Journal Article

**Welcome!**

To use the personalized features of this site, please **log in** or **register**.

If you have forgotten your username or password, we can **help**.

**My SpringerLink**

Marked Items

Alerts

Order History

**Saved Items**

All

Favorites



Effects of several 5'-carboxamide derivatives of adenosine on adenosine receptors of human platelets and rat fat cells

Journal Naunyn-Schmiedeberg's Archives of Pharmacology  
 Publisher Springer Berlin / Heidelberg  
 ISSN 0028-1298 (Print)  
 1432-1912 (Online)  
 Subject Biomedical and Life Sciences and Medicine  
 Issue Volume 327, Number 1 / March, 1984  
 DOI 10.1007/BF00504989  
 Pages 36-42  
 SpringerLink Sunday, December 12, 2004

**Add to marked items**

Add to shopping cart

Add to saved items

Recommend this article

**D. Ukena<sup>1</sup>, E. Böhme<sup>1</sup> and U. Schwabe<sup>1</sup>**

(1) Pharmakologisches Institut der Universität Heidelberg, Im Neuenheimer Feld 366, D-6900 Heidelberg 1, Federal Republic of Germany

**Received:** 13 February 1984 **Accepted:** 12 April 1984

**Summary** The effects of several 5'-carboxamide derivatives of adenosine on stimulatory ( $R_a$ ) adenosine receptors of human platelets and inhibitory ( $R_i$ ) adenosine receptors of rat fat cells have been compared. 5'-N-Cyclopropylcarboxamidoadenosine (CPCA) and 5'-N-ethylcarboxamidoadenosine (NECA) most potently inhibited ADP-induced aggregation of human platelets as shown by  $IC_{50}$ -values of 0.24 and 0.34

Find more options

 ... 

- ☒ Within this issue
- ☐ Within this journal
- ☐ Within all content

**Export this article**

Export this article as RIS | Text

**Text****PDF**

The size of this document is 820 kilobytes. Although it may be a lengthier download, this is the most authoritative online format.

Open: Entire document

**Referenced by**

2 newer articles

1. Cristalli, Gloria (2006) Highlights on the

$\mu\text{mol/l}$ . 5'-N-

Methylcarboxamidoadenosine (MECA;  $\text{IC}_{50}$  0.81  $\mu\text{mol/l}$ ) and 5'-N-carboxamidoadenosine (NCA;  $\text{IC}_{50}$  2.1  $\mu\text{mol/l}$ ) were less potent, whereas adenosine, 2-chloroadenosine and (-)  $\text{N}^6$ -phenylisopropyladenosine [(-)PIA] exhibit  $\text{IC}_{50}$ -values of about 1.5  $\mu\text{mol/l}$ . Nearly the same rank order of potency was obtained for stimulation of adenylate cyclase activity of platelet membranes and for inhibition of [ $^3\text{H}$ ] NECA binding to human platelets. In order to examine the effects of the carboxamide analogues on  $R_i$  adenosine receptors of rat fat cells inhibition of lipolysis and adenylate cyclase were studied. (-)PIA was the most potent inhibitor of lipolysis as shown by an  $\text{IC}_{50}$  of 0.5 nmol/l, followed by CPCA ( $\text{IC}_{50}$  1.1 nmol/l) and NECA ( $\text{IC}_{50}$  1.3 nmol/l), whereas MECA ( $\text{IC}_{50}$  17.9 nmol/l) and NCA ( $\text{IC}_{50}$  20.1 nmol/l) were much less potent than NECA in inhibiting lipolysis. Similar results were obtained for inhibition of adenylate cyclase activity of fat cell membranes and for competition with [ $^3\text{H}$ ]PIA binding to fat cell membranes. The relative potencies of the adenosine analogues at both receptor subclasses were calculated from the ratio of the  $\text{IC}_{50}$ -values for inhibition of platelet aggregation and of lipolysis. (-)PIA showed the highest selectivity for  $R_i$  receptors as indicated by a 2,900-fold lower  $\text{IC}_{50}$  for the antilipolytic than for the antiaggregatory effect. The  $R_a/R_i$  activity ratio for NECA was about 260, for CPCA 220, for NCA 105 and for MECA 45. These results indicate that all 5'-carboxamide adenosine derivatives are more potent agonists at  $R_i$  receptors than at  $R_a$  receptors. Since MECA has a higher

Development of A2A Adenosine Receptor Agonists and Antagonists. *ChemMedChem* [CrossRef]

- Gessi, S (2000) A2A adenosine receptors in human peripheral blood cells. *British Journal of Pharmacology* 129(1) [CrossRef]

[Ads by Google](#)

**MECA Electronics, Inc.**  
RF/Microwave Coaxial Components N SMA BNC 7/16 DIN TNC, DC-18 GHz  
[www.e-meca.com](http://www.e-meca.com)

**Adenosine Syringe Pump**

Advanced Infusion 12S Pro. Perfect for pharm stress.  
[cnmtstore.com](http://cnmtstore.com)

**Adenosine**

Get the information you need to make an informed choice.  
[www.dlife.com](http://www.dlife.com)

**Fix Meca.exe**

Download a Free Scan Tool & Fix Meca.exe Instantly!  
[www.TuneupAdvisor.com](http://www.TuneupAdvisor.com)

**Protein Characterization**

Protein Solution's DynaPro Now available only through Wyatt!  
[www.wyatt.com](http://www.wyatt.com)

[Advertise on this site](#)

*J Neurophysiol* 78: 1965-1972, 1997;  
0022-3077/97 \$5.00

The Journal of Neurophysiology Vol. 78 No. 4 October 1997, pp. 1965-1972  
Copyright ©1997 by the American Physiological Society

# NMDA-Independent LTP by Adenosine A<sub>2</sub> Receptor-Mediated Postsynaptic AMPA Potentiation in Hippocampus

Kofi Kessey<sup>1</sup> and David J. Mogul<sup>1, 2</sup>

<sup>1</sup> Department of Neurobiology and Physiology and <sup>2</sup> Department of Biomedical Engineering, Northwestern University, Evanston, Illinois 60208

## ► ABSTRACT

Kessey, Kofi and David J. Mogul. NMDA-independent LTP by adenosine A<sub>2</sub> receptor-mediated postsynaptic AMPA potentiation in hippocampus. *J. Neurophysiol.* 78: 1965-1972, 1997. The role of adenosine A<sub>2</sub> receptors in normal synaptic transmission

• Abstract  
▼ Introduction  
▼ Methods  
▼ Results  
▼ Discussion  
▼ References

and tetanus-induced long-term potentiation (LTP) was tested by stimulation of the Schaffer collateral pathway and recording of the field excitatory postsynaptic potential (EPSP) in the CA1 region of rat transverse hippocampal slices. Activation of adenosine A<sub>2</sub> receptors with the A<sub>2</sub> agonist N<sup>6</sup>-[2-(3,5-dimethoxyphenyl)-2-(2-methylphenyl)-ethyl]adenosine (DPMA; 20 nM) enhanced synaptic transmission during low-frequency test pulses (0.033 Hz). Paired stimulation before and during DPMA exposure indicated no paired-pulse facilitation as a result of A<sub>2</sub> activation, suggesting that enhancement was not a result of presynaptic modulation. DPMA enhanced the early phase  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) component of the EPSP. In contrast, DPMA had no effect on the N-methyl-D-aspartate (NMDA) component isolated using low extracellular Mg<sup>2+</sup> and the AMPA receptor blocker 6-cyano-7-nitroquinoxaline-2,3-dione (20  $\mu$ M), indicating that the effects of A<sub>2</sub> activation on synaptic transmission were mediated by a postsynaptic enhancement of the AMPA response. Activation of adenosine A<sub>2</sub> receptors during a brief tetanus (100 Hz, 1 s) increased the level of LTP by 36% over that seen in response to a tetanus under control conditions. DPMA exposure after prior induction of LTP showed no additional potentiation, indicating that the mechanisms that contribute to both types of increases in synaptic transmission share a common mechanism. A slow onset NMDA-independent LTP could be induced by application of a tetanus during perfusion of DPMA with the NMDA blocker AP5 (50  $\mu$ M).

### This Article

- [Abstract](#) FREE
- [Full Text \(PDF\)](#)
- [Alert me when this article is cited](#)
- [Alert me if a correction is posted](#)
- [Citation Map](#)

### Services

- [Email this article to a friend](#)
- [Similar articles in this journal](#)
- [Similar articles in PubMed](#)
- [Alert me to new issues of the journal](#)
- [Download to citation manager](#)
- [Cited by other online articles](#)

### Google Scholar

- [Articles by Kessey, K.](#)
- [Articles by Mogul, D. J.](#)
- [Articles citing this Article](#)

### PubMed

- [PubMed Citation](#)
- [Articles by Kessey, K.](#)
- [Articles by Mogul, D. J.](#)